


## Arylation and Vinylation of $\alpha$ -Diazocarbonyl Compounds with Boroxines

Cheng Peng, Wei Zhang, Guobing Yan, and Jianbo Wang

*Org. Lett.*, 2009, 11 (7), 1667-1670 • DOI: 10.1021/ol900362d • Publication Date (Web): 04 March 2009

Downloaded from <http://pubs.acs.org> on March 26, 2009



### More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)


Arylation and Vinylation of  
 $\alpha$ -Diazocarbonyl Compounds with  
BoroxinesCheng Peng,<sup>†</sup> Wei Zhang,<sup>†</sup> Guobing Yan,<sup>‡</sup> and Jianbo Wang<sup>\*†‡</sup>

Beijing National Laboratory of Molecular Sciences (BNLMS) and Key Laboratory of Bioorganic Chemistry and Molecular Engineering of Ministry of Education, College of Chemistry, Peking University, Beijing 100871, China, and Department of Chemistry, Tongji University, Shanghai 200092, China

wangjb@pku.edu.cn

Received February 19, 2009


## ABSTRACT



An alternative approach for  $\alpha$ -arylation and  $\alpha$ -vinylation of carbonyl compounds is described: reaction between aryl- or vinylboroxines with  $\alpha$ -diazocarbonyl compounds leads to the formation of  $\alpha$ -arylated or  $\alpha$ -vinylated carbonyl compounds under mild conditions.

$\alpha$ -Arylation and  $\alpha$ -vinylation of carbonyl compounds are very important in organic synthesis. Although various approaches have been developed for these purposes, they usually suffer drawbacks in one way or another. Recently, palladium-catalyzed  $\alpha$ -arylation and  $\alpha$ -vinylation of carbonyl compounds, which are mostly developed by Buchwald and Hartwig, have been shown to provide simple and efficient methods to introduce aryl and vinyl groups to the  $\alpha$  carbon of carbonyl compounds.<sup>1,2</sup> However, these catalytic processes have to be carried out under strong basic conditions. An

indirect approach toward  $\alpha$ -arylated carbonyl compounds would be the arylation of the corresponding  $\alpha$ -diazocarbonyl compounds, which could be easily available through diazotization of the corresponding carbonyl compounds under mild conditions (Scheme 1). In this paper, we report an alternative

Scheme 1.  $\alpha$ -Arylation of Carbonyl Compounds

$\alpha$ -arylation method by the reaction of  $\alpha$ -diazocarbonyl compounds with easily accessible arylboroxines. This approach also applies to  $\alpha$ -vinylation by using the corresponding vinylboroxines.

<sup>†</sup> Peking University.<sup>‡</sup> Tongji University.


(1) For seminal reports, see: (a) Palucki, M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1997**, *119*, 11108. (b) Hamann, B. C.; Hartwig, J. F. *J. Am. Chem. Soc.* **1997**, *119*, 12382. (c) Satoh, T.; Kawamura, Y.; Miura, M.; Nomura, M. *Angew. Chem., Int. Ed.* **1997**, *36*, 1740.

(2) For selected recent publications, see: (a) Hama, T.; Culkin, D. A.; Hartwig, J. F. *J. Am. Chem. Soc.* **2006**, *128*, 4976. (b) Takemiya, A.; Hartwig, J. F. *J. Am. Chem. Soc.* **2006**, *128*, 14800. (c) Martín, R.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2007**, *46*, 7236. (d) Huang, J.; Bunel, E.; Faul, M. M. *Org. Lett.* **2007**, *9*, 4343. (e) Liao, X.; Weng, Z.; Hartwig, J. F. *J. Am. Chem. Soc.* **2008**, *130*, 195. (f) Durbin, M. J.; Willis, M. C. *Org. Lett.* **2008**, *10*, 1413. (g) Vo, G. D.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2008**, *47*, 2127.

The reactions of  $\alpha$ -diazocarbonyl compound with organoboron compounds have been previously reported to form a C–C bond between alkyl or aryl and an  $\alpha$  carbon of carbonyl group.<sup>3–7</sup> Pioneering work by Hooz and co-workers established the alkylation of diazonitrile, ethyl diazoacetate, and diazoketones by means of trialkylboranes.<sup>3</sup> The reactions reported by Hooz are only limited to alkylboranes, and the reaction is sluggish with poor yields for organoboranes containing relatively bulky alkyl groups. Further improvement was made by Brown, Hooz, and co-workers by using dialkylchloroboranes and alkyl- or aryl-dichloroboranes, which are more reactive boron reagents toward diazo compounds.<sup>4</sup> The drawback of Brown's modification is that chlorinated byproducts are formed in some cases due to the competing 1,2-chloro migration. Particularly notable in this modification is that arylation can occur with high yields by using aryl-dichloroboranes,<sup>4b</sup> and most recently, Brown and Salunkhe have extended this reaction by using alkenyldichloroboranes.<sup>6</sup> However, these otherwise very attractive arylation and vinylation reactions have so far received limited attention, presumably because toxic and unstable boron compounds are used, and the reactions have to be carried out at low temperature. Moreover, in Brown's report, only ethyl diazoacetate is applied as a diazo substrate.

Mechanistically, it is reasonable to consider that these reactions are initiated by the nucleophilic attack of diazo substrate to boron. Thus, increasing the electronegativity in the boron should enhance the reaction. This explains the high reactivity with dialkylchloroboranes and alkydichloroboranes in Brown's reaction, as compared with the original trialkylboranes in Hooz's system. In view of the fact that organoboronic acids and their derivatives, which are stable, nontoxic, and easily available, are gaining popularity within the synthetic organic chemistry community,<sup>8,9</sup> we have conceived that it is worthwhile to investigate the reaction of diazo compounds with boronic acids and their derivatives.<sup>10</sup>

At the outset of this investigation, we examined the reaction of methyl  $\alpha$ -diazopropionate **1a** with phenylboronic acid **2** (eq 1). The  $\alpha$ -arylation product **3a** was isolated, albeit in low yield. Subsequent efforts to optimize this reaction proved fruitless. When phenyl pinacolborane **4** was used instead of boronic acid, **3a** was not observed, and most starting materials remained unchanged after stirring at 80 °C for 24 h. To our delight, when phenylboronic acid **2** was replaced by phenylboroxine **5a**, the reaction under the same conditions was found to be faster and the yield was slightly higher.



Further optimization of the reaction conditions was carried out. The results are summarized in Table 1. Changing solvent

**Table 1.** Optimization of Reaction Conditions with Methyl  $\alpha$ -Diazopropionate **1a** and Phenylboroxine **5a**<sup>a</sup>

| entry                 | additive (3 equiv)                  | solvent           | T (°C)    | yield <sup>b</sup> (%) |
|-----------------------|-------------------------------------|-------------------|-----------|------------------------|
| 1                     | non                                 | PhMe              | 80        | 50                     |
| 2                     | non                                 | PhMe              | 40        | 33                     |
| 3                     | non                                 | PhH               | 60        | 31                     |
| 4                     | non                                 | <i>n</i> -hexane  | 60        | 24                     |
| 5                     | non                                 | DCE               | 60        | 30                     |
| 6                     | Non                                 | Et <sub>3</sub> N | 60        | 38                     |
| 7                     | Et <sub>3</sub> N                   | PhMe              | 60        | 72                     |
| 8                     | pyridine                            | PhMe              | 60        | 68                     |
| 9                     | morpholine                          | PhMe              | 60        | 60                     |
| 10                    | <sup>i</sup> Pr <sub>2</sub> EtN    | PhMe              | 60        | 49                     |
| 11                    | <sup>i</sup> Pr <sub>2</sub> NH     | PhMe              | 60        | 82                     |
| 12                    | <sup>i</sup> Pr <sub>2</sub> NH     | DCE               | 60        | 88                     |
| <b>13<sup>c</sup></b> | <b><sup>i</sup>Pr<sub>2</sub>NH</b> | <b>DCE</b>        | <b>60</b> | <b>91</b>              |
| 14 <sup>c</sup>       | <sup>i</sup> Pr <sub>2</sub> NH     | THF               | 60        | 85                     |
| 15 <sup>d</sup>       | <sup>i</sup> Pr <sub>2</sub> NH     | DCE               | 60        | 79                     |

<sup>a</sup> If otherwise indicated, the reaction was carried out with 1.0 equiv of diazo compound, 1.5 equiv of (PhBO)<sub>3</sub>. <sup>b</sup> Yields calculated from GC using mesitylene as an internal standard. <sup>c</sup> 1.0 equiv of (PhBO)<sub>3</sub> was used. <sup>d</sup> 0.6 equiv of (PhBO)<sub>3</sub> was used.

and temperature seemed not improve the yields (entries 1–6). It was then observed that the reaction could be significantly improved by adding base as additives, among which diisopropylamine afforded the best results (entry 11). The reaction

(10) For a recent report on Pd-catalyzed coupling of arylboronic acids with  $\alpha$ -diazocarbonyl compounds, see: Peng, C.; Wang, Y.; Wang, J. *J. Am. Chem. Soc.* **2008**, *130*, 1566.

(3) (a) Hooz, J.; Linke, S. *J. Am. Chem. Soc.* **1968**, *90*, 5936. (b) Hooz, J.; Linke, S. *J. Am. Chem. Soc.* **1968**, *90*, 6891. (c) Hooz, J.; Morrison, G. F. *Can. J. Chem.* **1970**, *48*, 868. (d) Hooz, J.; Gunn, D. M. *J. Chem. Soc., Chem. Commun.* **1969**, 139. (e) Hooz, J.; Gunn, D. M. *Tetrahedron Lett.* **1969**, 3455. (f) Hooz, J.; Gunn, D. M. *J. Am. Chem. Soc.* **1969**, *91*, 6195.

(4) (a) Brown, H. C.; Midland, M. M.; Levy, A. B. *J. Am. Chem. Soc.* **1972**, *94*, 3662. (b) Hooz, J.; Bridson, J. N.; Calzada, J. G.; Brown, H. C.; Midland, M. M.; Levy, A. B. *J. Org. Chem.* **1973**, *38*, 2574.

(5) For further applications of the Hooz reaction, see: (a) Pasto, D. J.; Wojtkowski, P. W. *Tetrahedron Lett.* **1970**, 215. (b) Pasto, D. J.; Wojtkowski, P. W. *J. Org. Chem.* **1971**, *36*, 1790. (c) Newman, H. J. *J. Org. Chem.* **1974**, *39*, 100. (d) Hooz, J.; Bridson, J. N. *J. Am. Chem. Soc.* **1973**, *95*, 602. (e) Hooz, J.; Oudenes, J.; Roberts, J. L.; Benderly, A. *J. Org. Chem.* **1987**, *52*, 1347.

(6) Brown, H. C.; Salunkhe, A. M. *Synlett* **1991**, 684.

(7) For comprehensive reviews on diazo compounds, see: (a) Doyle, M. P.; McKervey, M. A.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*; Wiley-Interscience: New York, 1998. (b) Ye, T.; McKervey, M. A. *Chem. Rev.* **1994**, *94*, 1901. (c) Zhang, Z.; Wang, J. *Tetrahedron* **2008**, *64*, 6577.

(8) For reviews on Suzuki–Miyaura coupling, see: (a) Suzuki, A. *Acc. Chem. Res.* **1982**, *15*, 178. (b) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457. (c) Miyaura, N. *J. Organomet. Chem.* **2002**, *653*, 54. (d) Miyaura, N. In *Advances in Metal-Organic Chemistry*; Liebeskind, L. S., Ed.; JAI: London, 1998; Vol. 6, pp 187.

(9) For use of arylboroxines in organic synthesis, see: (a) Hayashi, T.; Senda, T.; Takaya, Y.; Ogasawara, M. *J. Am. Chem. Soc.* **1999**, *121*, 11591. (b) Chen, F.-X.; Kina, A.; Hayashi, T. *Org. Lett.* **2006**, *8*, 341.

could be further improved by using 1,2-dichloroethane (DCE) as solvent (entry 13). Since three phenyl groups in one phenylboroxine can be theoretically transferred to the product, we proceeded to reduce the amount of phenylboroxine. One equivalent of boroxine afforded satisfactory results (entry 13). When the phenylboroxine was further reduced, the reaction became sluggish and the yield was diminished due to the side reactions of diazo substrate (entry 15).

In Hooz and Brown's reports, only the diazo compounds without  $\alpha$ -substitution ( $N_2$ CHA, A = CN, CO<sub>2</sub>Et, COMe, CPh, etc.), in most cases the ethyl diazoacetate, were investigated. As a general  $\alpha$ -arylation method, more diazo substrates with structural diversity have to be employed. Thus, the optimized reaction conditions were applied to the reaction of **5a** with various  $\alpha$ -diazocarbonyl compounds (Table 2). The  $\alpha$ -phenylation products were obtained in

**Table 2.** Reaction of **1a–m** with Phenylboroxine **5a**<sup>a</sup>

| entry | diazosubstrate (1, R, R')                                    | time (h) | yield <sup>b</sup> (%) |
|-------|--|----------|------------------------|
| 1     | <b>1a</b> , Me, OMe  | 4        | <b>3a</b> , 84         |
| 2     | <b>1b</b> , <sup>n</sup> Pr, OMe                             | 9        | <b>3b</b> , 84         |
| 3     | <b>1c</b> , <sup>i</sup> Pr, OMe                             | 10       | <b>3c</b> , 65         |
| 4     | <b>1d</b> , Bn, OMe  | 12       | <b>3d</b> , 91         |
| 5     | <b>1e</b> , H, OEt   | 49       | <b>3f</b> , 90         |
| 6     | <b>1f</b> , H, N <sup>i</sup> Pr <sub>2</sub>                | 48       | <b>3e</b> , 96         |
| 7     | <b>1g</b> , Me, C <sub>6</sub> H <sub>5</sub>                | 43       | <b>3g</b> , 68         |
| 8     | <b>1h</b> , Me, <i>p</i> -ClC <sub>6</sub> H <sub>4</sub>    | 48       | <b>3h</b> , 61         |
| 9     | <b>1i</b> , Me, <i>p</i> -BrC <sub>6</sub> H <sub>4</sub>    | 48       | <b>3i</b> , 64         |
| 10    | <b>1j</b> , C <sub>6</sub> H <sub>5</sub> , OMe              | 16       | <b>3j</b> , 78         |
| 11    | <b>1k</b> , <i>p</i> -MeOC <sub>6</sub> H <sub>4</sub> , OMe | 2        | <b>3k</b> , 81         |
| 12    | <b>1l</b> , <i>p</i> -BrC <sub>6</sub> H <sub>4</sub> , OMe  | 2        | <b>3l</b> , 61         |
| 13    | <b>1m</b> , 1-naphthyl, OMe                                  | 17       | <b>3m</b> , 56         |

<sup>a</sup> The reaction was carried out with diazo compound (1.0 equiv) and phenylboroxine (1.0 equiv) at 60 °C (for entries 1–4, 6–10) or 100 °C (entries 5, 11–13). <sup>b</sup> Isolated yield.

moderate to high yields in all cases. Reaction with  $\alpha$ -alkyl substituted diazo esters gave the  $\alpha$ -phenyl-substituted aliphatic esters in good yields (entries 2–4). When the alkyl group was replaced by hydrogen, the reaction became sluggish but the yield remained high (entries 5). Reaction with  $\alpha$ -diazamide also worked well (entry 6), thus providing an efficient way for  $\alpha$ -arylation of amides.

The results for the reaction with  $\alpha$ -diazoketones are also summarized in Table 2 (entries 7–9). The slightly diminished yields in these cases as compared with  $\alpha$ -diazooesters might be attributed to the decreased reactivity of the substrates under the reaction conditions, which resulted in 1, 2-H shift side reactions. Finally, the  $\alpha$ -aryl- $\alpha$ -diazooacetates were also studied (entries 10–13). Reactions with these substrates needed a higher temperature to reach completion. The electron-donating group was found to facilitate the reaction

(entry 12). Since the  $\alpha$ -aryl- $\alpha$ -diazooacetates are easily available from the corresponding  $\alpha$ -arylacetates,<sup>11</sup> this process constitutes a practical access to  $\alpha, \alpha$ -diaryl-substituted acetates.

Furthermore, a series of arylboroxines were examined through the reaction with *p*-tolyl diazoacetate **6** under the same conditions at 100 °C. As shown in Table 3, the

**Table 3.** Reaction of Arylboroxine **5b–h** with **6**<sup>a</sup>


| entry          | <b>5</b> , Ar'   | time (h) | yield ( <b>7</b> , %) <sup>b</sup> |
|----------------|--|----------|------------------------------------|
| 1              | <b>5b</b> , <i>p</i> -CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>    | 13       | <b>7a</b> , 73                     |
| 2              | <b>5c</b> , <i>m</i> -CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>    | 14       | <b>7b</b> , 79                     |
| 3              | <b>5d</b> , <i>m</i> -ClC <sub>6</sub> H <sub>4</sub>                  | 14       | <b>7c</b> , 86                     |
| 4              | <b>5e</b> , <i>m, p</i> -Cl <sub>2</sub> C <sub>6</sub> H <sub>3</sub> | 4        | <b>7d</b> , 87                     |
| 5              | <b>5f</b> , <i>m</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>    | 2        | <b>7e</b> , 87                     |
| 6 <sup>c</sup> | <b>5g</b> , <i>m</i> -CHOC <sub>6</sub> H <sub>4</sub>                 | 5        | <b>7f</b> , 60                     |
| 7              | <b>5h</b> , <i>p</i> -CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>   | 13       | <b>7g</b> , 56                     |
| 8              | <b>5i</b> , 1-naphthyl   | 11       | <b>7h</b> , 56                     |

<sup>a</sup> Reaction conditions: **5b–i** (1.0 equiv), **6** (1.0 equiv). <sup>b</sup> Isolated yield. <sup>c</sup> *i*-Pr<sub>2</sub>NH was replaced by Et<sub>3</sub>N.

reactions all proceeded well to afford diaryl-substituted acetate products **7a–h**. It was noted that the reaction was significantly accelerated by electron-withdrawing substituents of arylboroxines (entries 4–6).

It is worthwhile to note that both the arylboroxine substrate and the diazo substrate in this reaction can bear halide substituents (Table 2, entries 8, 9, and 13; Table 3, entries 3 and 4). To introduce an aryl group bearing halide substituent to the  $\alpha$ -position of carbonyl group would be difficult with the Pd-catalyzed reaction due to the anticipated side reactions.

The scope of this  $\alpha$ -arylation approach was further explored by using cyclic  $\alpha$ -diazocarbonyl substrates (eqs 2 and 3). The reaction with cyclic  $\alpha$ -diazoketone **8** and cyclic  $\alpha$ -diazamide **10** under the same conditions provided the corresponding  $\alpha$ -arylation products in moderate yields.



Compared with  $\alpha$ -arylation,  $\alpha$ -vinylation of carbonyl compounds has only been sparsely reported.<sup>2d,12</sup> Encouraged by the success of the  $\alpha$ -arylation with arylboroxines, we turned our attention to investigate the corresponding  $\alpha$ -vi-

nylation by using vinylboroxine. With  $\alpha$ -diazo amide **1f**, vinylboroxines **12a–f** were applied to the similar reaction conditions, and the results are summarized in Table 4. In all

**Table 4.** Reaction of Vinylboroxines **12a–f** with **1f**<sup>a</sup>

| entry | <b>12</b> , R  | time (h) | yield ( <b>13</b> , %) <sup>b</sup> |
|-------|--|----------|-------------------------------------|
| 1     | <b>12a</b> , C <sub>6</sub> H <sub>5</sub>                           | 12       | <b>13a</b> , 65                     |
| 2     | <b>12b</b> , <i>p</i> -CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub> | 11       | <b>13b</b> , 88                     |
| 3     | <b>12c</b> , <i>p</i> -ClC <sub>6</sub> H <sub>4</sub>               | 10       | <b>13c</b> , 77                     |
| 4     | <b>12d</b> , 1-naphthyl  | 13       | <b>13d</b> , 74                     |
| 5     | <b>12e</b> , <i>n</i> -pentyl  | 24       | <b>13e</b> , 45                     |
| 6     | <b>12f</b> , BnOCH <sub>2</sub> CH <sub>2</sub> -                    | 22       | <b>13f</b> , 61                     |

<sup>a</sup> Reaction conditions: **1f** (1 equiv), **12a–f** (1 equiv). <sup>b</sup> Isolated yield.


cases, the reaction occurred smoothly, giving  $\alpha$ -vinylation products in moderate to good yields. The byproduct attributed from the shift of the double bond to the  $\alpha,\beta$  position were not detected in these cases. The same  $\alpha$ -vinylation reaction has also been tested with diazo ester **1a**. In that case, the major product of  $\beta,\gamma$ -unsaturated ester was accompanied by small amount of  $\alpha,\beta$  unsaturated ester as minor product.

The mechanism of this  $\alpha$ -arylation and  $\alpha$ -vinylation is thought to be similar to the reaction of  $\alpha$ -diazo compounds with alkylborane (Scheme 2).<sup>3,4</sup> The reaction is initiated by nucleophilic attack of the diazo carbon to boroxine, generating intermediate **A**, followed by 1,2-migration of R'' from

(11) (a) Davies, H. M. L.; Townsend, R. J. *J. Org. Chem.* **2001**, *66*, 6595. (b) Zhang, X.; Qu, Z.; Ma, Z.; Shi, W.; Jin, X.; Wang, J. *J. Org. Chem.* **2002**, *67*, 5621.

(12) For examples, see: (a) Ooi, T.; Goto, R.; Maruoka, K. *J. Am. Chem. Soc.* **2003**, *125*, 10494. (b) Kim, H.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2008**, *130*, 398.

**Scheme 2.** Mechanistic Rationale



boron to carbon to give **B**. Hydrolysis of **B** leads to the product. The additive of diisopropylamine has been found to significantly improve the yield of the reaction. It was observed that adding diisopropylamine could markedly slow down the reaction. Since diazo compounds are generally sensitive to acids, we speculate that the role of diisopropylamine is to neutralize the reaction system by interact with boroxine or acidic species generated during the reactions, thus preventing the diazo substrate from decomposition.

In summary, we have developed a general approach for  $\alpha$ -arylation and  $\alpha$ -vinylation by the reaction of  $\alpha$ -diazo carbonyl compounds with boroxines. Since the reaction conditions are mild and both diazo compounds and boroxines are easily available, this reaction should find an application in organic synthesis.

**Acknowledgment.** The project is generously supported by the Natural Science Foundation of China (Grant Nos. 20832002, 20772003, and 20821062), the National Basic Research Program of China (973 Program, No. 2009CB825300), and the MOE of China (Cheung Kong Scholars Program).

**Supporting Information Available:** Experiment procedure, characterization data, <sup>1</sup>H and <sup>13</sup>C NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL900362D